


AD-A265 501



DTIC  
ELECTE  
JUN 09 1993  
S A D

## PUBLICATION REPORT

This document has been approved  
for public release and sale; its  
distribution is unlimited.

1739

17/93

COMBINED MICROAUTORADIOGRAPHIC AND HISTOPATHOLOGIC ANALYSIS  
OF THE FATE OF CHALLENGE SCHISTOSOMA MANSONI SCHISTOSOMULA  
IN MICE IMMUNIZED WITH IRRADIATED CERCARIAE

BY

Olakunle O. Kassim, David A. Dean, Beverly L. Mangold  
and Franz von Lichtenberg

U.S. NAVAL MEDICAL RESEARCH UNIT NO. 3  
(CAIRO, ARAB REPUBLIC OF EGYPT)  
PSC 452, BOX 5000  
FPO AE 09835-0007

93-12885



93 6 08 083

## COMBINED MICROAUTORADIOGRAPHIC AND HISTOPATHOLOGIC ANALYSIS OF THE FATE OF CHALLENGE *SCHISTOSOMA MANSONI* SCHISTOSOMULA IN MICE IMMUNIZED WITH IRRADIATED CERCARIAE

OLAKUNLE O. KASSIM, DAVID A. DEAN, BEVERLY  
L. MANGOLD, AND FRANZ VON LICHTENBERG

Department of Microbiology, Howard University College of Medicine, Washington, DC; U. S. Naval  
Medical Research Unit No. 3, Cairo, Egypt; Department of Pathology, Harvard Medical School and  
Brigham and Women's Hospital, Boston, Massachusetts

**Abstract.** Combined microautoradiographic and histopathologic methods were used to locate and examine schistosomula of *Schistosoma mansoni* in the lungs of irradiated cercaria-immunized mice 21 days after percutaneous challenge infection with <sup>75</sup>Se-labeled cercariae. Of 75 schistosomula examined in serial sections, 53% were located in the pulmonary microvasculature, 23% in alveolar spaces, 3% with one end in a vessel and the other in an alveolar space, and the locations of 21% were not identified. Inflammatory reactions of variable intensity were observed around schistosomula in both vascular and alveolar sites, although the most intense category of reactions was associated almost entirely with alveolar larvae. All autoradiographic foci contained recognizable schistosomula. Although the concentration of reduced silver grains precluded cytostructural analysis, observations on schistosomular contour and shape provided no evidence of larval damage. Our findings suggest that immune elimination of schistosomula in mice immunized with irradiated cercariae is partly or largely effected by a process of alveolar extrusion of viable parasites during their lung migration.

Evidence from histologic and autoradiographic studies has indicated that the lungs represent a major site of *Schistosoma mansoni* elimination in both naive and immunized hosts.<sup>1-11</sup> The mechanisms responsible for this elimination and for the delay in migration to and from the lungs seen in most immune models have not been clearly defined. During the period when larvae disappear from the body, intense inflammatory foci, possibly representing sites previously occupied by larvae, have been observed in the lungs by light microscopy.<sup>6</sup> These foci were significantly more numerous in mice immunized with irradiated cercariae than in controls.

Over the last 70 years, there have been numerous observations of schistosomula in pulmonary air spaces.<sup>7, 12-19</sup> Most recently, Crabtree and Wilson used an ingenious method to identify ultrathin sections of lung tissue containing radioactivity associated with <sup>75</sup>Se-labeled larvae.<sup>7</sup> Examination of these sections by electron microscopy revealed that in both normal mice and mice immunized with irradiated cercariae, schistosomula shifted from vascular to alveolar sites with increased time spent in the lungs. Dead or

damaged schistosomula were not detected in either site.

In the present study, we have combined microautoradiographic and histopathologic techniques to examine by light microscopy the autoradiographic foci produced in the lungs by <sup>75</sup>Se-labeled schistosomula. This approach complements those previously used to search for and examine schistosomula undergoing elimination. Conventional light microscopic images obtained with thin sections<sup>4</sup> and free of autoradiographic grains<sup>4, 6</sup> provide a more detailed picture of schistosomular morphology, but the addition of autoradiography provides a means of searching for sites occupied by larvae that have disintegrated beyond the point of recognition. The autoradiographic-electron microscopic method<sup>7</sup> provides a better means of assessing the condition of individual schistosomula, particularly those showing early or subtle damage, while light microscopy makes it possible to examine more larvae and to examine them in complete sets of serial sections.

The results of this microscopic study, coupled with macroautoradiographic tracking data for the




FIGURE 1. Schistosomulum of *Schistosoma mansoni* migrating along an interalveolar capillary, cut at its full length. Note the absence of an inflammatory reaction (hematoxylin and eosin stained, original magnification  $\times 360$ ).



FIGURE 2. A somewhat contracted schistosomulum of *Schistosoma mansoni* occupying a dilated pulmonary venule. Note the absence of an inflammatory reaction (hematoxylin and eosin stained, original magnification  $\times 360$ ).

same batch of immunized mice,<sup>11</sup> our previous macroautoradiographic studies,<sup>1,2</sup> and other light and electron microscopic observation,<sup>4,6,7</sup> indicate that new hypotheses are needed to explain the process of schistosome elimination from the lungs.

#### MATERIALS AND METHODS

##### Host and parasite

The female C57Bl/6J mice used in this study were obtained from the Jackson Laboratories (Bar Harbor, ME). A Puerto Rican strain of *S. mansoni* maintained in *Biomphalaria glabrata* was used.<sup>20</sup> The cercariae used for immunization were exposed to 50 kilorads of gamma radiation from a cesium-137 source at a rate of 1,300 rads/min. Mice were immunized by three percutaneous (tail immersion) exposures to 539–572 irradiated cercariae at 0, 101, and 129 days. Twenty-six days after the third immunization, two mice were challenged by tail immersion exposure to 1,243 cercariae (mean of 1,228 penetrants) labeled with <sup>75</sup>Se-L-selenomethionine as previously described.<sup>1</sup> Both mice were killed 21 days after challenge. (Immunity of this batch of immunized mice to challenge infection was demonstrated in

the accompanying report,<sup>11</sup> using the same pools of immunizing and challenge cercariae.)

Lung-stage schistosomula were obtained by exposure of mice to 3,000–4,000 radiolabeled cercariae and recovery of schistosomula seven days later by mincing and incubation of lungs.<sup>11</sup> The schistosomula were killed by suspending them in 1.0 ml of Earle's lactalbumin hydrolyzate medium (Gibco, Grand Island, NY) containing 5% normal mouse serum in a 12-ml glass conical tube and immersing the tube in a swirling 50°C bath for 5 min. They were then diluted with the same medium and approximately 2,100 schistosomula in 0.4 ml were introduced into each of two immunized mouse by tail vein injection.

##### Histologic and macroautoradiographic analysis

Twenty-one days after percutaneous challenge or two days after intravenous injection of killed lung schistosomula, mice were killed by inhalation of CO<sub>2</sub>. The lungs were slowly inflated by tracheal injection with neutral 10% formalin, stored in the same formalin, and embedded in paraffin. A portion of the lungs of each mouse was serially cut into 10- $\mu$  sections. All sections were dipped in Eastman Kodak (Rochester, NY)



FIGURE 3. Comma-shaped schistosomulum of *Schistosoma mansoni* in an alveolar space, outside the border of an adjacent, dense inflammatory focus (hematoxylin and eosin stained, original magnification  $\times 360$ ).



FIGURE 4. Shortened, broadened schistosomulum of *Schistosoma mansoni* in an atrial pulmonary air sac adjacent to an inflammatory focus (above) and to a distended pulmonary capillary (below) (hematoxylin and eosin stained, original magnification  $\times 360$ ).

NTB emulsion, air-dried, and stored in light-excluding black boxes at 4°C for 12–14 weeks. The slides were then developed in D-19 developer (Eastman Kodak) for 3 min, washed twice in distilled water, fixed for 2 min in acetic acid, washed for 10 min in distilled water, and lightly stained with hematoxylin and eosin.

Cellular reactions to schistosomula were classified as negative, with no focal accumulation of inflammatory cells above background levels; + (mild), with leukocyte density in the vicinity of schistosomula somewhat above background levels; ++ (moderate), with dense focal leukocytic reactions 2–5-cells thick or more diffuse reactions intermediate between + and +++; +++ (intense), with large, dense reactions more than five cells thick, which were usually highly focal and completely enveloping larvae and/or surrounding alveoli.

## RESULTS

### Percutaneous challenge

Immunized mice in this experiment were resistant to a challenge infection with 155 of the same batch of cercariae used for microautoradiography; adult worm burdens were reduced 75.3% relative to challenge controls.<sup>11</sup>

Autoradiographic foci were examined in 10- $\mu$  sections of the lungs of two immunized mice 21 days after challenge with <sup>75</sup>Se-labeled cercariae. Every autoradiographic focus found contained a schistosomulum. No schistosomula free of autoradiographic foci were found. Eighty-one serial sections (39 continuous sections from one mouse and 42 from the second) were examined. Seventy-five schistosomula and associated autoradiographic foci were examined in complete sets of serial sections (35 from one mouse and 40 from the other).

Fifty-three percent of the schistosomula observed were clearly within blood vessels (Figures 1 and 2), and 23% were clearly in air spaces (Figures 3 and 4). Three percent had one end in a vessel and the other end in an air space. The locations of the remaining 21% could not be determined (Table 1). A few alveolar larvae were associated with hemorrhages rather than with inflammatory foci.

Intense (+++) inflammatory responses were associated with 42.1% of the alveolar schistosomula, compared with only one (2.5%) intravascular schistosomulum (Table 2). Approximately twice as many vascular schistosomula were associated with mild (+) or moderate (++) reactions as were alveolar schistosomula, and 37.5% of vascular schistosomula and 26.3% of

TABLE 1

*Distribution of Schistosoma mansoni schistosomula in the lungs of irradiated cercaria-immunized mice 21 days after challenge with <sup>75</sup>Se-labeled cercariae\**

Location of schistosomula	No. (%)
Blood vessels	40 (53)
Alveolar spaces	17 (23)
Both†	2 (3)
Undetermined location	16 (21)

\* The mice examined were from experiment no. 4 reported in the accompanying paper.<sup>11</sup>

† These two schistosomula clearly had one end in a vessel and the other end in an alveolar space. Several in the undetermined location category were possibly of this type.

alveolar schistosomula were completely free of focal inflammation. Most cellular reactions consisted of small and large mononuclear cells, together with some neutrophils and eosinophils. In some cases, granulocytes outnumbered mononuclear cells, while a few reactions were composed almost entirely of small mononuclear cells with granulocytes limited to their perimeters, resembling granulomas.

Brush-like inflammatory reactions,<sup>6</sup> either distal to or in the vicinity of schistosomula, were often seen in ++ reactions, especially those associated with vascular schistosomula. The +++ reactions associated with alveolar schistosomula were all dense and highly focal. In some cases, the cellular infiltrates filled the air spaces, obscuring their contours and surrounding the schistosomula. In others, the cells were limited to the adjacent tissue, revealing intact schistosomula free within the air spaces.

Both alveolar and vascular schistosomula appeared to be free of structural damage. Even when surrounded by intense inflammatory foci, no attachment of leukocytes to schistosomular surfaces could be detected. However, subtle parasite damage or cell attachment could not be conclusively ruled out by histologic analysis, since the thickness of the tissue sections and the density of autoradiographic foci often restricted detailed visualization.

#### *Comparison of heat-killed with live challenge schistosomula*

Two days after intravenous injection of heat-killed <sup>75</sup>Se-labeled lung schistosomula into the lung vasculature of immunized mice, all schistosomula had disintegrated beyond recognition. Nevertheless, numerous dense inflammatory foci

TABLE 2

*Intensity of inflammatory cellular responses associated with Schistosoma mansoni schistosomula in the lungs of irradiated cercaria-immunized mice 21 days after challenge infection with <sup>75</sup>Se-labeled cercariae*

Inflammatory response	% of schistosomula		
	Vascular	Alveolar	Undetermined
N (none)	37.5	26.3	18.8
+ (mild)	15.0	10.5*	6.2
++ (moderate)	45.0	21.1	18.8
+++ (intense)	2.5	42.1*	56.2

\* Includes one schistosomulum with one end in a blood vessel.

were seen, and all of these foci contained schistosomular debris with overlying silver grains, identifiable as autoradiographic foci. Conversely, all autoradiographic foci observed were associated with dense inflammatory reactions. Although the autoradiographic foci produced by lung squashes were macroscopically found to be less numerous and less intense on day 2 than on the day of injection,<sup>11</sup> it is likely that the microautoradiographic method was sensitive enough to allow detection of injected larvae on day 2. Following a percutaneous challenge, numerous focal inflammatory reactions free of recognizable schistosomula were observed in the lungs, as had been reported previously.<sup>6</sup> None of these reactions was associated with concentrations of reduced silver grains that might represent the remains of a disintegrating schistosomulum.

#### DISCUSSION

Under most experimental conditions, the loss of *S. mansoni* from hosts immunized with irradiated cercariae following a challenge infection can be largely accounted for by the disappearance of schistosomula during the lung phase of migration.<sup>4-10, 22</sup> This was confirmed in the present study by macroautoradiographic observations in the accompanying report.<sup>11</sup> Exceptions that have been reported are elimination in the skin of monkeys,<sup>23</sup> elimination in the skin of CBA/Ca mice immunized with the Mill Hill strain of *S. mansoni*,<sup>24-27</sup> and additional larval elimination in the liver of guinea pigs.<sup>23, 28</sup>

von Lichtenberg and others observed that parasite-free inflammatory foci after challenge were significantly more numerous in the lungs of immunized mice than in control mice, and they

suggested that such residual inflammatory foci might represent sites of worm elimination.<sup>6</sup> At that time, testing of this hypothesis was precluded by the lack of markers for detecting schistosomular secretions or remnants in inflammatory foci. In the present study, <sup>75</sup>Se-L-selenomethionine provided such a marker. However, our examination failed to detect any parasite material in the lungs apart from that associated with structurally intact, apparently undamaged schistosomula. Clusters of reduced silver grains in the absence of larvae were seen only in sections adjacent to those containing larvae, and these sections were frequently completely free of silver grains. Similarly, Crabtree and Wilson did not find evidence of significant structural damage to lung-stage schistosomula of *S. mansoni* in their study.<sup>7</sup>

An important question is whether or not radioactive material left behind by disintegrating larvae would persist long enough in focal concentrations to allow detection of killing sites by autoradiography. In this study, heat-killed radiolabeled schistosomula injected into the lung vasculature continued to produce detectable autoradiographic foci for several days after the larvae had disintegrated beyond recognition. Similarly, larvae that died in the skin<sup>29</sup> and the lungs<sup>4</sup> from the effects of prior irradiation are associated with dense inflammatory foci prior to death, but can be identified morphologically (and therefore presumably by autoradiography) for some time after disintegration has begun. Therefore, it seems likely that any larvae that died in pulmonary inflammatory foci would leave behind radioactive evidence.

An alternative interpretation of parasite-free inflammatory lung foci is that they represent sites previously occupied by schistosomula, but subsequently vacated by them. This is an attractive hypothesis, but is difficult to test. As discussed above, radiolabeled larval products do not extend far beyond the schistosomular surface and therefore probably do not provide a reliable means of locating sites recently vacated by intact larvae.

The results of this study confirm the electron microscopic observations of Crabtree and Wilson,<sup>7</sup> which indicate that most *S. mansoni* schistosomula present in the lungs of irradiated cercaria-immunized mice after percutaneous challenge infection are intact and free of attached host leukocytes, in spite of the fact that they are

often surrounded by large numbers of inflammatory cells. The finding that the most intense inflammatory reactions were associated almost exclusively with schistosomula located in alveoli (Table 2) could be due to the triggering of inflammatory responses by host tissue damage resulting from breaching of the capillary-alveolar barrier, the washing away of schistosomular exoantigens from schistosomula located in blood vessels, or both factors.

In conclusion, we believe there are three ways in which schistosomula could be eliminated from the lungs. First, schistosomula are destroyed in the lungs, but <sup>75</sup>Se-labeled parasite and host material is removed from the killing sites too quickly to be detectable as discrete autoradiographic foci. This seems unlikely in view of the present microscopic observations on migrating larvae, coupled with the data from previous studies on the numbers of parasites that disappear from the host lungs and body during the second and third weeks after challenge,<sup>3,11</sup> and the fact that killed injected larvae and dying irradiated larvae<sup>4,29</sup> are histologically detectable for several days. Second, the disappearing schistosomula may actively migrate out of the lungs by the vascular route and subsequently be destroyed in post-lung migration sites. Evidence against this proposal is provided by the failure of previous macroautoradiographic tracking studies to detect a post-lung site of larval accumulation other than the liver.<sup>3,11</sup> Also, a great effort would presumably be required for schistosomula to return to blood vessels from alveolar spaces,<sup>10</sup> especially if surrounded by inflammatory cells. Third, intact live schistosomula may be eliminated from alveoli via the upper airways, and subsequently, via the gastrointestinal tract. This possibility was first suggested by Donato-Cioli and has been proposed by Crabtree and Wilson.<sup>7</sup> Evidence for such a mechanism has been provided by autoradiographic detection of schistosomula in the trachea and in the lumina of the esophagus, stomach, and intestines of nonimmune mice.<sup>30</sup> Additional evidence based on qualitative analysis of macroautoradiographic foci is presented in the accompanying report.<sup>11</sup>

**Acknowledgment:** The experiments reported herein were conducted according to the principles set forth in the current edition of the *Guide for the Care and Use of Laboratory Animals*, Institute of Laboratory Animal Resources, National Research Council.

Financial support: This work was supported by the Naval Medical Research and Development Command Work Units no. 3M161102.BS10.AF.426 and 3M161102.BS13.AK.311, and the Office of Naval Research Contract no. N00014-81-0552.

Disclaimer: The opinions or assertions contained herein are the private ones of the authors and are not to be construed as official or reflecting the views of the U. S. Navy Department, Department of Defense, or the U. S. Government.

Authors' addresses: Olakunle O. Kassim, Department of Microbiology, Howard University College of Medicine, Washington, DC 20059. David A. Dean and Beverly L. Mangold, U. S. Naval Medical Research Unit No. 3, PSC 452, FPO AE 09835-0007. Franz von Lichtenberg, Department of Pathology, Harvard Medical School and Brigham and Women's Hospital, Boston, MA 02115.

Reprint requests: Research Publications, U. S. Naval Medical Research Unit No. 3, PSC 452 Box 5000, FPO AE 09835-0007.

#### REFERENCES

1. Dean DA, Mangold BL, Georgi JR, Cioli D, 1982. Studies on the induction and expression of resistance to schistosome infection in mice. *Pontificiae Academiae Scientiarum Scripta Varia* 47: 73-77.
2. Georgi JR, Dean DA, Mangold BL, 1983. Schistosoma mansoni: temporal distribution of radio-selenium-labelled schistosomula in lungs of mice during the first two weeks of infection. *Parasitology* 86: 31-36.
3. Mangold BL, Dean DA, 1983. Autoradiographic analysis of Schistosoma mansoni migration from skin to lungs in naive mice. *Am J Trop Med Hyg* 32: 785-789.
4. Mastin AJ, Bickle QD, Wilson RA, 1983. Schistosoma mansoni: migration and attrition of irradiated and challenge schistosomula in the mouse. *Parasitology* 87: 87-102.
5. Dean DA, Mangold BL, Georgi JR, Jacobson RH, 1984. Comparison of Schistosoma mansoni migration patterns in normal and irradiated cercaria-immunized mice by means of autoradiographic analysis. *Am J Trop Med Hyg* 33: 89-96.
6. von Lichtenberg F, Correa-Oliveira R, Sher A, 1985. The fate of challenge schistosomula in the murine anti-schistosome vaccine model. *Am J Trop Med Hyg* 34: 96-106.
7. Crabtree JE, Wilson RA, 1986. The role of pulmonary cellular reactions in the resistance of vaccinated mice to Schistosoma mansoni. *Parasite Immunol* 8: 265-286.
8. Knopf PM, Cioli D, Mangold BL, Dean DA, 1986. Migration of Schistosoma mansoni in normal and passively immunized laboratory rats. *Am J Trop Med Hyg* 35: 1173-1184.
9. Wilson RA, Coulson PS, Dixon B, 1986. Migration of the schistosomula of Schistosoma mansoni in mice vaccinated with radiation-attenuated cercariae, and normal mice: an attempt to identify the timing and site of parasite death. *Parasitology* 92: 101-116.
10. Coulson PS, Wilson RA, 1982. Examination of the mechanisms of pulmonary phase resistance to Schistosoma mansoni in vaccinated mice. *Am Trop Med Hyg* 38: 529-539.
11. Dean DA, Mangold BL, 1992. Evidence that both normal and immune elimination of Schistosoma mansoni take place at the lung stage of migration prior to parasite death. *Am J Trop Med Hyg* 47: 238-248.
12. Miyagawa Y, Takemoto S, 1921. The mode of infection of Schistosomum japonicum and principal route of its journey from the skin to the portal vein in the host. *J Pathol Bacteriol* 24: 168-174.
13. Koppisch E, 1937. Studies on schistosomiasis mansoni in Puerto Rico. IV. The pathological anatomy of experimental schistosomiasis mansoni in the rabbit and albino rat. *Puerto Rico J Public Health Trop Med* 13: 1-54.
14. Kagan IG, Meranze DR, 1958. The histopathology of experimental infections in mice with Schistosomatium douthitti. *Am J Trop Med Hyg* 7: 285-293.
15. Sadun EH, Lin SS, Williams JE, 1958. Studies on the host parasite relationships to Schistosoma japonicum. I. The effect of single graded infections and the route of migration of schistosomula. *Am J Trop Med Hyg* 7: 494-499.
16. Lin SS, Sadun EH, 1959. Studies on the host parasite relationships to Schistosoma japonicum. V. Reactions in the skin, lungs and liver of normal and immune animals following infection with Schistosoma japonicum. *J Parasitol* 45: 549-556.
17. Magalhaes-Filho A, 1959. Pulmonary lesions in mice experimentally infected with Schistosoma mansoni. *Am J Trop Med Hyg* 8: 527-535.
18. von Lichtenberg F, Ritchie LS, 1961. Cellular resistance against schistosomula of Schistosoma mansoni in Macaca mulatta monkeys following prolonged infections. *Am J Trop Med Hyg* 10: 859-869.
19. Wilks NE, 1967. Lung-to-liver migration of schistosomes in the laboratory mouse. *Am J Trop Med Hyg* 16: 599-605.
20. Stirewalt MA, Uy A, 1969. Schistosoma mansoni: cercarial penetration and schistosomule collection in an in vitro system. *Exp Parasitol* 26: 17-28.
21. Dean DA, Minard P, Murrell KD, Vannier WE, 1978. Resistance of mice to secondary infection with Schistosoma mansoni. II. Evidence for a correlation between egg deposition and worm elimination. *Am J Trop Med Hyg* 27: 957-965.
22. Ford MJ, Bickle QD, Taylor MG, Andrews BJ, 1984. Passive transfer of resistance and the site of immune-dependent elimination of the challenge infection in rats vaccinated with highly

- irradiated cercariae of *Schistosoma mansoni*. *Parasitology* 89: 461-482.
23. Hsu SYL, Hsu HF, Penick GD, Lust GL, Osborne JW, 1974. Dermal hypersensitivity to schistosome cercariae in rhesus monkeys during immunization and challenge. I. Complex hypersensitivity reactions of a well-immunized monkey during the challenge. *J Allergy Clin Immunol* 54: 339-349.
  24. Miller KL, Smithers SR, 1980. *Schistosoma mansoni*: the attrition of a challenge in mice immunized with highly irradiated live cercariae. *Exp Parasitol* 50: 212-221.
  25. McLaren DJ, Pearce EJ, Smithers SR, 1985. Site potential for challenge attrition in mice, rats and guinea pigs vaccinated with irradiated cercariae of *Schistosoma mansoni*. *Parasite Immunol* 7: 29-44.
  26. Kamiya H, Smithers SR, McLaren DJ, 1987. *Schistosoma mansoni*: autoradiographic tracking studies of isotopically-labeled challenge parasites in naive and vaccinated CBA/Ca mice. *Parasite Immunol* 9: 515-529.
  27. McLaren DJ, Strath M, Smithers SR, 1987. *Schistosoma mansoni*: evidence that immunity in vaccinated and chronically infected CBA/Ca mice is sensitive to treatment with a monoclonal antibody that depletes cutaneous effector cells. *Parasite Immunol* 9: 667-682.
  28. McLaren DJ, Rogers MV, 1986. *Schistosoma mansoni*: liver phase challenge attrition is a stage-dependent phenomenon in guinea-pigs vaccinated with highly irradiated cercariae. *Parasite Immunol* 8: 307-318.
  29. Byram JE, von Lichtenberg F, Lewis FA, Stirewalt MA, 1983. Pathology of a live attenuated anti-schistosome vaccine in mice. *Am J Trop Med Hyg* 32: 94-105.
  30. Georgi JR, Wade SE, Dean DA, 1987. *Schistosoma mansoni*: mechanisms of attrition and routes of migration from lungs to hepatic portal system in the laboratory mouse. *J Parasitol* 73: 706-711.

Accession For	
NTIS CRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	20



REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Service, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Project Director, Paperwork Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE 1992	3. REPORT TYPE AND DATES COVERED		
4. TITLE AND SUBTITLE Combined Microautoradiographic and Histopathologic Analysis of the Fate of Challenge <u>Schistosoma mansoni</u> Schistosomula in Mice Immunized with Irradiated Cercariae		5. FUNDING NUMBERS PE- 61102A WU- 3M161102BS13.AK 311		
6. AUTHOR(S) Kassim, Olakunle O., Dean, David A., Mangold, Beverly L. and von Lichtenberg, Franz				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) U.S. Naval Medical Research Unit No. 3 PSC 452, Box 5000 FPO AE 09835-0007		8. PERFORMING ORGANIZATION REPORT NUMBER 17/93		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Naval Medical Research and Development Command, National Naval Medical Center Building 1, Tower 12 Bethesda, MD 20889-5044		10. SPONSORING/MONITORING AGENCY REPORT NUMBER		
11. SUPPLEMENTARY NOTES Published in: Am. J. Trop. Med. Hyg., 47(2):231-237, 1992; Acc. No. 1739.				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; Distribution is unlimited.		12b. DISTRIBUTION CODE		
13. ABSTRACT (Maximum 200 words) Combined microautoradiographic and histopathologic methods were used to locate and examine schistosomula of <u>Schistosoma mansoni</u> in the lungs of irradiated cercaria-immunized mice 21 days after percutaneous challenge infection with <sup>75</sup> Se-labeled cercariae. Of 75 schistosomula examined in serial sections, 53% were located in the pulmonary microvasculature, 23% in alveolar spaces, 3% with one end in a vessel and the other in an alveolar space, and the locations of 21% were not identified. Inflammatory reactions of variable intensity were observed around schistosomula in both vascular and alveolar sites, although the most intense category of reactions was associated almost entirely with alveolar larvae. All autoradiographic foci contained recognizable schistosomula. Although the concentration of reduced silver grains precluded cytostructural analysis, observations on schistosomular contour and shape provided no evidence of larval damage. Our findings suggest that immune elimination of schistosomula in mice immunized with irradiated cercariae is partly or largely effected by a process of alveolar extrusion of viable parasites during their lung migration.				
14. SUBJECT TERMS <u>Schistosoma mansoni</u> ; Schistosomula challenge; Microautoradiographic and histopathologic analysis; Irradiated cercariae; Mice.			15. NUMBER OF PAGES 8	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT	